



Illness Policy – Service Users

When groups of people are together, illness and disease can spread from one person to another even when the recommended hygiene and infection control practices have been implemented. A policy on illness establishes a shared expectation of IOE's exclusion guidelines.

IOE is committed to preventing the spread of infectious illnesses through the implementation, monitoring and maintaining of simple strategies such as, handwashing, effective cleaning procedures, and an understanding and knowledge of the health of people who access the service.

In the case of any illness or infectious disease, due consideration should be given to the following:

- a. To avoid the spread of infection to anyone else at IOE
- b. To recover properly
- c. To minimise disruption to the workplace

The parent or responsible person must contact the relevant IOE co-ordinator, or the IOE after hours emergency phone if out of normal office hours, when the person attending the service is ill. The person must not attend the OSHC program in order to protect the health and safety of others and should remain away in line with the exclusion periods in Figure 1

Refer the cancellation policy for information regarding cancellation fees and notice periods.

All illness that are highly infectious and cause an outbreak need to be reported to the Department of Health and Department of Education.

Figure 1 – Minimum Exclusion Periods for Common Infectious Diseases

Number	Conditions	Exclusion of cases	Exclusion of contacts
1	Chickenpox	Exclude until all blisters have dried. This is usually at least 5 days after the rash appears in unimmunised children, but may be less in previously immunised children	Any child with an immune deficiency (for example, leukaemia) or receiving chemotherapy should be excluded for their own protection. Otherwise not excluded
2	Conjunctivitis	Exclude until discharge from eyes has ceased	Not excluded
3	Cytomegalovirus (CMV) infection	Exclusion is not necessary	Not excluded
4	Diarrhoeal illness	Exclude until there has not been vomiting or a loose bowel motion for 24 hours	Not excluded
5	Diphtheria	Exclude until medical certificate of recovery is received following at least two negative throat swabs, the first not less than 24 hours after finishing a course of antibiotics and the other 48 hours later	Exclude family/household contacts until cleared to return by the Chief Health Officer
6	Glandular fever (Epstein-Barr Virus infection)	Exclusion is not necessary	Not excluded
7	Hand, Foot and Mouth disease	Exclude until all blisters have dried	Not excluded
8	Haemophilus influenzae type b (Hib)	Exclude until 48 hours after initiation of effective therapy	Not excluded
9	Hepatitis A	Exclude until a medical certificate of recovery is received, but not before 7 days after the onset of jaundice or illness	Not excluded
10	Hepatitis B	Exclusion is not necessary	Not excluded
11	Hepatitis C	Exclusion is not necessary	Not excluded
12	Herpes (cold sores)	Young children unable to comply with good hygiene practices should be excluded while the lesion is weeping. Lesions to be covered by dressing, where possible	Not excluded
13	Human immunodeficiency virus infection (HIV)	Exclusion is not necessary	Not excluded
14	Impetigo	Exclude until appropriate treatment has commenced. Sores on exposed surfaces must be covered with a watertight dressing	Not excluded
15	Influenza and influenza like illnesses	Exclude until well	Not excluded unless considered necessary by the Chief Health Officer
16	Leprosy	Exclude until approval to return has been given by the Chief Health Officer	Not excluded
17	Measles	Exclude for at least 4 days after onset of rash	Immunised contacts not excluded. Unimmunised contacts should be excluded until 14 days after the first day of appearance of rash in the last case. If unimmunised contacts are vaccinated within 72 hours of exposure with any infectious case, or received Normal Human Immunoglobulin (NHIG) within 144 hours of exposure of any infectious case, they may return to the facility
18	Meningitis (bacterial other than meningococcal meningitis)	Exclude until well	Not excluded
19	Meningococcal infection	Exclude until adequate carrier eradication therapy has been completed	Not excluded if receiving carrier eradication therapy
20	Mumps	Exclude for 5 days or until swelling goes down (whichever is sooner)	Not excluded
21	Molluscum contagiosum	Exclusion is not necessary	Not excluded
22	Pertussis (Whooping cough)	Exclude the child for 21 days after the onset of cough or until they have completed 5 days of a course of antibiotic treatment	Contacts aged less than 7 years in the same room as the case who have not received three effective doses of pertussis vaccine should be excluded for 14 days after the last exposure to the infectious case, or until they have taken 5 days of a course of effective antibiotic treatment
23	Poliovirus infection	Exclude for at least 14 days from onset. Re admit after receiving medical certificate of recovery	Not excluded
24	Ringworm, scabies, pediculosis (head lice)	Exclude until the day after appropriate treatment has commenced	Not excluded
25	Rubella (German measles)	Exclude until fully recovered or for at least four days after the onset of rash	Not excluded
26	Severe Acute Respiratory Syndrome (SARS)	Exclude until medical certificate of recovery is produced	Not excluded unless considered necessary by the Chief Health Officer
27	Shiga toxin or Verotoxin producing Escherichia coli (STEC or VTEC)	Exclude if required by the Chief Health Officer and only for the period specified by the Chief Health Officer	Not excluded
28	Streptococcal infection (including scarlet fever)	Exclude until the child has received antibiotic treatment for at least 24 hours and the child feels well	Not excluded
29	Tuberculosis (excluding latent tuberculosis)	Exclude until receipt of a medical certificate from the treating physician stating that the child is not considered to be infectious	Not excluded
30	Typhoid fever (including paratyphoid fever)	Exclude until approval to return has been given by the Chief Health Officer	Not excluded unless considered necessary by the Chief Health Officer